INTERNAL MEDICINE www.drshavindergill.com 2415 Ware Street, Abbotsford, BC V2S 3C6 TEL (604) 870-3277

<u>(It is a Tudor style home with 2 wrought iron gates; look for yellow polls</u>. It is beside <u>Mill Lake Park on the same side of the street</u>) <u>Please inform all referred patients to us of our office policies</u>

- 1. They <u>MUST</u> confirm with us, one week before their appointment. If they don't confirm with our office, then we will cancel their appointment <u>without</u> <u>any exceptions</u>. If a message is left no return call will be given.
- 2. If they do not show up for their appointment, confirmed or not, they will be sent a bill for the missed appointment. The cost will be \$100.
- 3. If they want to cancel their appointment, it must be cancelled 24 hours before the appointment time or there will be full charge for the missed appointment.
- 4. All patients should bring all their medications with them, including herbal and nonprescription medication.
- 5. <u>If you are pregnant or there is any possibility of pregnancy, then skin testing</u> <u>can not be done. Please inform us if you are pregnant.</u>
- 6. Skin prick testing will be done on your forearm. Please wear loose comfortable clothing.

- a. NO ANTIHISTAMINES, such as, Benadryl (Diphenhydramine), Dimetapp allergy, Gravol, Optimine, Periactin, Dramamine, Phenergan Tavist, Tylenol cold/allergy should be discontinued 48 hours before allergy testing
- b. Aerius(desloratadine), Allegra (Fexofenadine), Claritin (Loratadine), Atarax (Hydroxyzine), Reactine (Cetirizine), Blexten (Bilastine), Rupall (Rupatadine) and Zyrtec should be discontinued 72 hours before allergy testing.
- c. Elavil (Amitriptyline), Anafranil (Clomipramine), Sinequan (Doexepin), Tofranil (Imipramine), Surmontil (Trimipramine), Norpramin (Desipramine), Aventyl (Nortiplyline), and Vivaactil should be discontinued 7 days before allergy testing
- d. Celexa, Effexor, Prozac, Luvox, Paxil, Zoloft, Wellbutrin, Zyban, Serzone, and Desryl (Trazodone) should be discontinued 7 days before allergy testing.
- e. NO HEARTBURN MEDICATIONS, such as, Axid, Cimetidine, Pepcid and Zantac should be discontinued 48 hours before allergy testing.
- f. No Singulair (Montelukast) on the day of allergy testing.

INTERNAL MEDICINE www.drshavindergill.com 2415 Ware Street, Abbotsford, BC V2S 3C6 TEL (604) 870-3277

<u>(It is a Tudor style home with 2 wrought iron gates; look for yellow polls</u>. It is beside <u>Mill Lake Park on the same side of the street</u>) <u>Please inform all referred patients to us of our office policies</u>

- 1. They <u>MUST</u> confirm with us, one week before their appointment. If they don't confirm with our office, then we will cancel their appointment <u>without</u> <u>any exceptions</u>. If a message is left no return call will be given.
- 2. If they do not show up for their appointment, confirmed or not, they will be sent a bill for the missed appointment. The cost will be \$100.
- 3. If they want to cancel their appointment, it must be cancelled 24 hours before the appointment time or there will be full charge for the missed appointment.
- 4. All patients should bring all their medications with them, including herbal and nonprescription medication.
- 5. <u>If you are pregnant or there is any possibility of pregnancy, then skin testing</u> <u>can not be done. Please inform us if you are pregnant.</u>
- 6. Skin prick testing will be done on your forearm. Please wear loose comfortable clothing.

- a. NO ANTIHISTAMINES, such as, Benadryl (Diphenhydramine), Dimetapp allergy, Gravol, Optimine, Periactin, Dramamine, Phenergan Tavist, Tylenol cold/allergy should be discontinued 48 hours before allergy testing
- b. Aerius(desloratadine), Allegra (Fexofenadine), Claritin (Loratadine), Atarax (Hydroxyzine), Reactine (Cetirizine), Blexten (Bilastine), Rupall (Rupatadine) and Zyrtec should be discontinued 72 hours before allergy testing.
- c. Elavil (Amitriptyline), Anafranil (Clomipramine), Sinequan (Doexepin), Tofranil (Imipramine), Surmontil (Trimipramine), Norpramin (Desipramine), Aventyl (Nortiplyline), and Vivaactil should be discontinued 7 days before allergy testing
- d. Celexa, Effexor, Prozac, Luvox, Paxil, Zoloft, Wellbutrin, Zyban, Serzone, and Desryl (Trazodone) should be discontinued 7 days before allergy testing.
- e. NO HEARTBURN MEDICATIONS, such as, Axid, Cimetidine, Pepcid and Zantac should be discontinued 48 hours before allergy testing.
- f. No Singulair (Montelukast) on the day of allergy testing.

INTERNAL MEDICINE www.drshavindergill.com 2415 Ware Street, Abbotsford, BC V2S 3C6 TEL (604) 870-3277

<u>(It is a Tudor style home with 2 wrought iron gates; look for yellow polls</u>. It is beside <u>Mill Lake Park on the same side of the street</u>) <u>Please inform all referred patients to us of our office policies</u>

- 1. They <u>MUST</u> confirm with us, one week before their appointment. If they don't confirm with our office, then we will cancel their appointment <u>without</u> <u>any exceptions</u>. If a message is left no return call will be given.
- 2. If they do not show up for their appointment, confirmed or not, they will be sent a bill for the missed appointment. The cost will be \$100.
- 3. If they want to cancel their appointment, it must be cancelled 24 hours before the appointment time or there will be full charge for the missed appointment.
- 4. All patients should bring all their medications with them, including herbal and nonprescription medication.
- 5. <u>If you are pregnant or there is any possibility of pregnancy, then skin testing</u> <u>can not be done. Please inform us if you are pregnant.</u>
- 6. Skin prick testing will be done on your forearm. Please wear loose comfortable clothing.

- a. NO ANTIHISTAMINES, such as, Benadryl (Diphenhydramine), Dimetapp allergy, Gravol, Optimine, Periactin, Dramamine, Phenergan Tavist, Tylenol cold/allergy should be discontinued 48 hours before allergy testing
- b. Aerius(desloratadine), Allegra (Fexofenadine), Claritin (Loratadine), Atarax (Hydroxyzine), Reactine (Cetirizine), Blexten (Bilastine), Rupall (Rupatadine) and Zyrtec should be discontinued 72 hours before allergy testing.
- c. Elavil (Amitriptyline), Anafranil (Clomipramine), Sinequan (Doexepin), Tofranil (Imipramine), Surmontil (Trimipramine), Norpramin (Desipramine), Aventyl (Nortiplyline), and Vivaactil should be discontinued 7 days before allergy testing
- d. Celexa, Effexor, Prozac, Luvox, Paxil, Zoloft, Wellbutrin, Zyban, Serzone, and Desryl (Trazodone) should be discontinued 7 days before allergy testing.
- e. NO HEARTBURN MEDICATIONS, such as, Axid, Cimetidine, Pepcid and Zantac should be discontinued 48 hours before allergy testing.
- f. No Singulair (Montelukast) on the day of allergy testing.

INTERNAL MEDICINE www.drshavindergill.com 2415 Ware Street, Abbotsford, BC V2S 3C6 TEL (604) 870-3277

<u>(It is a Tudor style home with 2 wrought iron gates; look for yellow polls</u>. It is beside <u>Mill Lake Park on the same side of the street</u>) <u>Please inform all referred patients to us of our office policies</u>

- 1. They <u>MUST</u> confirm with us, one week before their appointment. If they don't confirm with our office, then we will cancel their appointment <u>without</u> <u>any exceptions</u>. If a message is left no return call will be given.
- 2. If they do not show up for their appointment, confirmed or not, they will be sent a bill for the missed appointment. The cost will be \$100.
- 3. If they want to cancel their appointment, it must be cancelled 24 hours before the appointment time or there will be full charge for the missed appointment.
- 4. All patients should bring all their medications with them, including herbal and nonprescription medication.
- 5. <u>If you are pregnant or there is any possibility of pregnancy, then skin testing</u> <u>can not be done. Please inform us if you are pregnant.</u>
- 6. Skin prick testing will be done on your forearm. Please wear loose comfortable clothing.

- a. NO ANTIHISTAMINES, such as, Benadryl (Diphenhydramine), Dimetapp allergy, Gravol, Optimine, Periactin, Dramamine, Phenergan Tavist, Tylenol cold/allergy should be discontinued 48 hours before allergy testing
- b. Aerius(desloratadine), Allegra (Fexofenadine), Claritin (Loratadine), Atarax (Hydroxyzine), Reactine (Cetirizine), Blexten (Bilastine), Rupall (Rupatadine) and Zyrtec should be discontinued 72 hours before allergy testing.
- c. Elavil (Amitriptyline), Anafranil (Clomipramine), Sinequan (Doexepin), Tofranil (Imipramine), Surmontil (Trimipramine), Norpramin (Desipramine), Aventyl (Nortiplyline), and Vivaactil should be discontinued 7 days before allergy testing
- d. Celexa, Effexor, Prozac, Luvox, Paxil, Zoloft, Wellbutrin, Zyban, Serzone, and Desryl (Trazodone) should be discontinued 7 days before allergy testing.
- e. NO HEARTBURN MEDICATIONS, such as, Axid, Cimetidine, Pepcid and Zantac should be discontinued 48 hours before allergy testing.
- f. No Singulair (Montelukast) on the day of allergy testing.